

Sleep, Health, and Human Capital: Evidence from Daylight Saving Time

Lawrence Jin*

Nicolas R. Ziebarth**

April 2019

Abstract

Chronic sleep deprivation is a significant and understudied public health issue. Using BRFSS survey data from the United States and an administrative census of 160 million hospital admissions from Germany, we study the causal relationship between sleep and health. Our empirical approach exploits the end of Daylight Saving Time in a quasi-experimental setting on a daily basis. First, we show that setting clocks back by one hour in the middle of the night significantly extends people's sleep duration. Second, we show that this nighttime extension reduces the share of people who involuntarily fall asleep during the day. In addition, we find significant health benefits via sharp reductions in hospital admissions. For example, hospitalizations due to cardiovascular diseases decrease by ten per day, per one million population. Using an event study approach, we find that the effects persists for four days after the time shift. Admissions due to heart attacks and injuries also exhibit the same characteristic four-day decrease. We provide a series of checks to rule out alternative mechanisms, and show that increasing sleep produces the human capital improvements. Finally, we discuss the benefits of additional sleep for the sleep-deprived as well as policy implications for nudging people to sleep more. Our findings illustrate the importance of public policies that target sleep deprivation.

Keywords: sleep deprivation, health, human capital, hospital admissions, BRFSS, Daylight Saving Time (DST)

JEL codes: H41, I18, I31

We thank Christian Bunnings, Ben Cowen, Peter Eibich, Osea Giuntella, Mike Grossman, Tatiana Homonoff, Ted Joyce, Bob Kaestner, Don Kenkel, Dean Lillard, Helmut Lüdtkepohl, Sara Markowitz, Joe Newhouse, Frank Schilbach, Hendrik Schmitz, Neeraj Sood, Shinsuke Tanaka, Gert Wagner, Hendrik Wolff as well as participants at the 2019 NBER Health Economics Spring Meeting, the Winter Meeting of the Econometric Society in San Francisco, the Health Economics and Policy Seminar at the University of Oxford, the Health Economics, Health Behaviors and Disparities Seminar at Cornell University, the University of Paderborn, and the GC Winter Workshop at DIW Berlin for very helpful comments and discussions. A special thank goes to Aline Paßlack for excellent research assistance. We take responsibility for all remaining errors in and shortcomings of the article.

*National University of Singapore, Centre for Behavioural Economics, 3 Research Link #02-01, Singapore 117602, e-mail: ljin@nus.edu.sg, Phone: +65 8177-1232

**Corresponding author: Cornell University, Policy Analysis and Management (PAM), 426 Kennedy Hall, Ithaca, NY 14853, DIW Berlin, and IZA Bonn, e-mail: nrz2@cornell.edu, Phone: +1-(607) 255-1180, Fax: +1-(607) 255-4071.

1. INTRODUCTION

Sleep deprivation has become a public health epidemic in recent years (CDC 2014). One third of the U.S. population report that they usually sleep less than the recommended minimum of 7 hours per night (Ford et al. 2015; Liu et al. 2016, Sheehan et al. 2019). The Centers for Disease Control and Prevention (CDC) warn that insufficient sleep leads to greater risk of car accidents and work injuries, as well as many chronic diseases and conditions such as high blood pressure, coronary heart disease, stroke, mental distress, and all-cause mortality.

Although the economic consequences of sleep deprivation are possibly substantial (Hillman et al. 2006; Mullainathan 2014), sleep has received relatively little attention in the economics literature until recently. Biddle and Hamermesh (1990) show that increases in time in the labor market reduce sleep. Sleep can also be impacted by television schedules (Hamermesh, 2008) and access to high-speed internet (Billari et al., 2018). Moreover, recent studies have identified causal relationships between inadequate sleep and reduced cognitive performance (Carrell et al. 2011; Giuntella et al. 2017), reduced wage returns (Gibson and Schrader, 2018), higher car accidents (Smith 2016), and higher incidences of obesity and diabetes (Giuntella and Mazzona, 2019). Hillman et al. (2006) estimate the economic costs of sleeplessness at almost one percent of GDP.

In this paper, we investigate whether increasing people's sleep reduces hospital admissions. To do this, we exploit the quasi-experimental nature of a regulation that affects the sleep pattern of more than one billion people in 70 countries around the globe: Daylight Saving Time (DST). It is the practice of setting clocks forward by one hour in spring and backward by one hour in the fall. Today, all countries in the European Union, the great majority of the U.S. states and Canadian provinces, as well as 40 other countries such as Mexico, Chile, Israel, and Iran observe DST.

Our identification strategy focuses on the time shift *in the fall* when the clocks “fall back” in the middle of the night; this regulation extends the duration of the night by one hour. We hypothesize that the additional hour allows especially sleep-deprived people to sleep more and find consistent evidence using a large U.S. survey. We find a significant increase in self-reported sleep duration following the “fall back”, and a significant decrease in the share of people who report having unintentionally fallen asleep during the day.

Next, we use the German Hospital Census to estimate the impact of the “fall back” on hospital admission rates across several disease categories. Exploiting all 160 million hospitalizations that occurred in Germany between 2000 and 2008 allows us to comprehensively control for seasonal and weekday confounders while maintaining enough statistical power to precisely identify population health effects at a daily level. We estimate changes in outcomes at the daily level compared to the neighboring weeks before and after the time shift while netting out seasonal and day-of-week effects.

Our findings show significant, sharp decreases in hospital admissions after the nighttime extension. Hospitalizations due to cardiovascular diseases decrease by ten per one million population, per day. This decrease lasts for four days. We find similar results across several disease categories and they are robust to multiple specifications. Our findings are in line with a large strand of the medical literature that has documented adverse physiological consequences of sleep restrictions (e.g. Moore et. al., 2002; Taheri et al., 2004; Berk et al., 2008; Mullington, et al., 2009; Killgore, 2010; Spaeth et al., 2013; for a review, see Banks and Dinges, 2007). We corroborate our main findings with permutation tests using non-DST transition weeks during the year. We also conduct falsification tests using outcomes that have no theoretical link with sleep, such as receiving a flu shot in the previous year. Moreover, we discuss alternative mechanisms through which the

DST transition might affect health, such as the shift in daylight, and discuss why these mechanisms are unlikely to explain our findings. In the last part of the paper, we monetize the economic benefits of increasing sleep at the population level.

This paper contributes to the human capital literature in economics. Since the seminal contributions by Becker (1964), Grossman (1972) and more recently by Heckman (e.g. Cunha and Heckman, 2007), many studies have theoretically modeled and empirically tested for human capital effects. Health is central component of human capital. For instance, studies have tested for the short and long-run effects of risky health behaviors (Cawley and Ruhm, 2011), ambient air pollution (Graff Zivin and Neidell, 2013; Currie et al., 2014), early life shocks (Kesternich et al. 2015) or *in utero* conditions (Almond and Currie, 2011).

This paper also relates to studies that have utilized daylight saving time as an empirical strategy. However, the large majority of DST studies have focused on spring DST. They have shown that falling back in time affects crime rates (Doleac and Sanders, 2015), traffic accidents (Hicks et al., 1998; Smith, 2016), energy demand (Kotchen and Grant, 2011; Sexton and Beatty, 2014), as well as our well-being (Kountouris and Remoundou, 2004; Kuehnle and Wunder, 2016). A large number of medical and psychology studies have also investigated the relationship between daylight saving time and sleep (e.g. Van Dongen et al, 2003; Lahti et al, 2008; Barnes and Wagner, 2009; Janszky et al. 2012; Jiddou et al. 2013; for a review see Harrison, 2013).

The next section briefly describes the data. Section 3 outlines the empirical methodology. Section 4 presents and discusses the findings and Section 5 concludes.

2. DATASETS

We employ a two-step approach in our analyses. First, we use a large U.S. survey to test if people sleep more when the night extends by one hour through the fall DST transition. Second, we utilize administrative hospital data from Germany to test for the impact of increased sleep on hospitalizations across various disease categories.

2.1 The U.S. Behavioral Risk Factor Surveillance System (BRFSS)

The BRFSS is a large annual telephone survey of U.S. adults aged 18 and above, which is administered by the CDC. The survey began in 1984 with fifteen participating states; by 1996, all 51 U.S. states participated in the survey. It is, by design, representative of state populations. In 2009, several states have started to include questions on sleep duration in the survey; this question expanded to all states between 2013 and 2016.

We focus on this period from 2013 to 2016, which includes 1.9 million survey responses in total. As shown in Figure 1, we extract six weeks around the time shift to ensure responses at a similar time of the year, to consider seasonalities in response behavior. Doing that, we obtain 174,503 survey responses in the main sample. Further, we include a robust set of time controls in our analysis, including month and day-of-week fixed effects.

Dependent Variables

The question on sleep duration reads: “*On average, how many hours of sleep do you get in a 24-hour period? Think about the time you actually spend sleeping or napping, not just the amount of sleep you think you should get.*” The answers are integers between 0 and 24. People on average report 7 hours of sleep, with a standard deviation of 1.5 (Table A1, Appendix). 32% report having slept 6 or fewer hours, which suggests a high level of sleep deprivation in the U.S.

However, the sleep question does not explicitly ask for the duration of sleep *last night*. Given the phrasing “24-hour period” and the emphasis on thinking about the time actually spent sleeping, the answers will likely be a weighted average of subjects’ sleep duration in the very recent past, with significant weight given to the previous night’s sleep. This measurement error will downward bias our estimates, as the true effect of the DST transition on sleep will only partially be reflected in the subjects’ responses. In other words, our estimates will likely be a lower bound of the true sleep effect.

We supplement the estimation with another survey question about tiredness during the day. Between 2009 and 2010,¹ the BRFSS asked: “*During the past 30 days, for about how many days did you find yourself unintentionally falling asleep during the day?*” The responses are integers between 0 and 30. We convert them into a binary variable of whether the subject has unintentionally fallen asleep in the past 30 days. On average, 35% of people report having unintentionally fallen asleep in the past 30 days (Table A1, Appendix).

If the fall nighttime extension induces people to sleep more, we should see an increase in the sleep duration and a similar decrease in people falling unintentionally asleep during the day. We discuss issues related to measurement errors of both variables in Appendix B.

Daylight Saving Time in the United States

In the United States, DST ends on the first Sunday in November. The time change occurs at 2am, where the clocks are set back to 1am, effectively extending the night by one hour. DST is observed by most states. Our empirical strategy only uses states that observe DST.

¹ This survey question was only asked during these two years.

2.2 German Hospital Admissions Census

The second dataset provides objective health measures. The dataset comprises all German hospital admissions from 2000 to 2008. The 16 German states collect these information and the *German Federal Statistical Office* provides restricted data access for researchers. Germany has about 82 million inhabitants and about 17 million hospital admission per year. To obtain the working dataset, we aggregate the admission-level data on the daily county level and then normalize admissions per 100,000 population. The data include information on age and gender, the day of admission, the county of residence as well as the diagnosis in form of the ICD-10 code.

As with the BRFSS, our working dataset focuses on the six weeks around the time shift (Figure 1). This main sample has 336,604 county-day observations.² We leave the data at the county-level and do not further aggregate up to the national level for a few reasons. This allows us to stratify the effects by county characteristics. Another reason is that we lose statistical power when aggregating up to a time series at the national level.

Dependent Variables

First, we generate *all cause admission rate*. On a given day, we observe 59.77 hospital admissions per 100,000 population (Table A1, Appendix).

Next, by extracting the ICD-10 codes I00-I99, we generate *cardiovascular admission rate*, the single most important subgroup of admissions (9.53 admissions per 100,000 population, Table A1). Extracting the codes I20 and I21, the *heart attack rate* is 1.59 admissions per 100,000 population.

² Between 2000 and 2008, Germany had up to 468 different counties. Mostly, due to mergers and reforms of the administrative boundaries, the number of counties varies across years.

Finally, we generate the *injury rate* (V01-X59) as well as the *respiratory* (J00-J99), *metabolic* (E00-E90), *neoplastic* (C00-D48), and *infectious admission rate* (A00-B99). We also test for changes in *drug overdosing* (T40) per 1 million population.

Daylight Saving Time in Germany

In Germany, DST ends on the last Sunday of October in all German states. The time change occurs on 3am where the clocks are set back to 2am.

3. EMPIRICAL SPECIFICATION

Our identification strategy relies on a plausibly exogenous extension of night sleep created by the nighttime extension through the end of DST in the fall. The transitions occur on different dates each year. Our large datasets allow us to comprehensively control for seasonal confounders, weekday effects, and yet still precisely estimate the health effects. Our preferred empirical specification identifies the effects at the daily level. We also estimate models at the weekly level to capture medium-term and potential intertemporal substitution effects.

3.1 Main Specification

Our preferred specification employs daily dummies around the DST time shift in the fall:

$$y_{id} = \beta_0 + \beta_1 DST_d + Vacation_d + DOW * \phi_m + \phi_m * \delta_t + X_{id} \gamma + \mu_s + \varepsilon_{id} \quad (1)$$

Where y_{id} is the outcome variable. For example, using the German Hospital Census it stands for admission rates in county i on day d . DST is a vector of fifteen daily dummies around the end of DST, $-7, -6, \dots, 0, \dots, 6, 7$, where 0 indicates the day of the time shift.

Equation (1) includes controls that net out seasonal and weekday confounders. These are crucial when using high-frequency data within the DST context. For example, hospital admissions

decrease on Sundays and on national holidays (Witte et al., 2005). *Vacation*_{*id*} controls for public holidays and Halloween.³

Due to the relevance of day-of-week (DOW) effects, we additionally interact DOW with month fixed effects ($DOW*\phi_m$). This is important, as Sundays in November may be systematically different from Sundays in September. For example, in our data, relative to Sundays, hospital admissions almost double on Mondays and this effect varies over the months of a year. Because DST ends always on Sundays, it is crucial to net out DOW effects by month of the year.

Our model also routinely includes month-year fixed effects ($\phi_m*\delta_t$) and some specifications additionally include linear and quadratic time trends at the annual level. However, the findings are robust to replacing month-year fixed effects with separate month and year fixed effects and omitting time trends. In addition, Equation (1) corrects for county-level or individual-level socio-demographics ($X_{id}'\gamma$) and persistent differences across states or counties (μ_s).

Because it is unlikely that county-level admission rates are either independent over time or across space, we correct the standard errors, ε_{id} , by applying two-way clustering across counties and over time (Cameron et al., 2011). When using the independently drawn and representative observations of the cross-sectional BRFSS, we cluster standard errors only at the date level (as it is no panel). All BRFSS regressions are probability weighted.

³ In Germany, official school vacations vary at the level of the 16 states by date, and also in lengths. Fall vacations lie between the beginning of October and mid-November, and vary by state, both in term of time and length. In the U.S., we include a dummy for Halloween, which occurs on October 31st each year. Halloween is only a very recent phenomenon in Germany. However, the German findings are robust to including Halloween fixed effects.

3.2 Identification

The key idea of our identification strategy is that DST transitions create plausibly exogenous variations in people’s sleep duration by extending the nighttime by one hour. Because fall DST simply extends night sleep for those who want to sleep more, we argue that it is a relatively clean setting without severe confounding factors (unlike in spring where the media regularly warns about drastic health effects and urges vulnerable people to take action).⁴

Turning the clocks back in the middle of the night is arguably exogenous to individuals. Our main specification de-trends the outcome variables using day-of-week by month and month-year fixed effects, in addition to the other controls in Equation (1). We also disentangle weekday and seasonal effects from vacation days or national holidays. The richness of our data still allows us to obtain precise estimates at the daily level. However, we also compare the day-to-day short-term effect of the change in time to the net effect on a weekly basis. Moreover, in effect heterogeneity specifications that test for behavioral mechanisms, we stratify the results by ambient climatic conditions such as temperatures and hours of sunshine.

Sample Selection

As illustrated by Figure 1, we restrict our main sample to three weeks before and three weeks after the time shift. Our preferred specification focuses on the week after DST as treatment week

⁴ Our findings are broadly consistent with the medical and psychology literature on DST transitions and sleep, although these studies overwhelmingly focus on spring and employ different empirical strategies (cf. Lahti et al, 2008; Barnes and Wagner, 2009; Janszky et al. 2012; Jiddou et al. 2013; Harrison 2013). Interestingly, as shown in Figure A2 for injuries, although we find single daily significant increases in admissions on the Monday and Tuesday after spring DST, we do not find spring effects that are as clean and clear as the fall effects (cf. Jin and Ziebarth, 2016). We hypothesize that this is a result of possible behavioral adjustments in spring. For example, to the extent that vulnerable people follow the very salient medical advice in the media, they likely adjust their bedtime schedules to ensure that they sleep enough, cf. “Your Daylight Saving Time Survival Guide” (Van Hare, 2019) or “It’s Daylight Saving Time! 6 Tips to Help You Deal with the Change” (Carroll, 2016).

and presents the main findings in an event study type graph by plotting 15 daily dummy estimates $-7, -6, \dots, 0, \dots, 6, 7$, where 0 represents the time shift. In other words, after netting out day-of-week, seasonal and other controls, Equation (1) compares the daily outcomes in the week before and after fall DST to four additional control weeks, as shown by Figure 1.

To check if the results are sensitive to this six-week sample selection around DST, we also estimate the models using all 52 weeks of the year, and the results remain robust. The findings are also robust to assigning all three post-transition weeks to the “treatment group.” Doing this yields results that are similar to a standard Regression Discontinuity design (cf. Hausman and Rapson, 2018), where the post-treatment outcomes are compared to that of the pre-treatment, conditional on all covariates shown in Equation (1), see for example Doleac and Sanders (2015).⁵

[Insert Figure 1 about here]

4. RESULTS

4.1 The Effects of Fall Back on Sleep Duration

Using the BRFSS, we first estimate the impact of the nighttime extension on sleep duration and tiredness. Without correcting for any seasonalities or other background characteristics, Figure 2 simply plots the mean reported sleep at the daily level for (a) those who were interviewed just before and after the DST transition, and (b) those who were interviewed in the weeks before and after. The black vertical line represents the DST transition that occurs for the dark blue line. As seen, the sleep duration for (b) remains relatively stable across all seven days of the week. Moreover, the sleep duration for (a) exhibits a trend that is very similar on Thursday to Saturday

⁵ The results are available upon request.

just before the transition. However, reported sleep then sharply increases on the Sunday of the nighttime extension, and this increase lasts for three days. In particular, the difference on Monday is statistically significant at the 1% level. On Tuesday, it is significant at the 5% level.

[Insert Figures 2 and 3 about here]

Next, we estimate a regression model as in Equation (1) and test if the results are robust to the inclusion of seasonal and other controls. Figure 3a plots the regression coefficients of the fifteen daily dummies in Equation (1), with *hours of sleep* as dependent variable. The *x*-axis represents the days relative to the nighttime extension (0 is the Sunday of the transition), and the *y*-axis shows the effect on sleep duration. Again, we see a sharp increase in self-reported sleep on the Monday following the transition. This effect persists for several days and is consistent with Figure 2. Together, they provide consistent evidence that people sleep more when clocks fall back in fall in the middle of the night.

We further corroborate these findings with a few more tests. Figure 3b plots the daily dummies of Equation (1) with *unintentionally fell asleep* as dependent variable. Consistent with our priors, we observe a sharp decline in the share of people who unintentionally fell asleep during the day after the nighttime extension. The decline is immediate and largest on the Sunday of the transition; and it persists for about four days before dissipating. The relatively clear pattern is even more reassuring when put into context because the measure is again a weighted average of the participants' reported tiredness in the past 30 days, creating a downward bias in our estimates. The sample is also relatively small due to the limited data availability. Despite these challenges, we find a significant decrease in tiredness at the daily level following the nighttime extension.

[Insert Table 1 about here]

Finally, Table 1 estimates the effect on sleep for the entire week of the time shift. That is, we run Equation (1) but replace the daily *DST* dummies with a binary *Week of Transition* indicator that equals one for the entire week of the transition (from Sunday of the transition until the Saturday after). According to column (1), *on average*, people sleep an additional 0.026 hours per night for seven days, for a total gain of 11 minutes of sleep throughout the week. This estimate is statistically significant at the 1% level. Note that this is a *population average*, which is also averaged over the entire DST week. Figures 2 and 3 suggest that the effect is concentrated during the first days after the transition. Moreover, the effect is very likely also concentrated among certain population subgroups, that is, the sleep deprived. As a comparison, Giuntella and Mazzonna (2019) find that a permanent sleep difference of 19 minutes leads to significantly higher obesity and blood pressure rates.

Column (2) turns to our measure of tiredness during the day. People are, on average, 4.4 percentage points less likely to unintentionally fall asleep during the treatment week, which is equivalent to a 13% decrease. The estimated weekly effect is statistically significant at the 5% level.

To summarize, while our sleep measures are self-reported, the results consistently provide evidence in support of the idea that the nighttime extension effectively extends people's sleep. Our sleep variables are likely downward-biased, but we still find effects, even at the daily level.

4.2 The Effects of Fall Back on Hospital Admissions

Next, using a census of hospital admissions for Germany from 2000 to 2008, we investigate whether the nighttime extension had any effect on hospital admissions. Table 2 shows the estimates by disease groups per 100,000 population in Germany. Each column is one model as in Equation (1) but the main regressor of interest is a dummy indicating the week of DST transition.

[Insert Table 2 about here]

Except for drug overdosing, all estimates are negative and highly significant, mostly at the 1% level. The weekly decreases in daily admissions range from 8.3% for the *all cause admission rate* (column (1)) to a similar 7.5% for *cardiovascular admissions* (column (2)). *Injuries* decrease by almost 5% or about 2.7 per 1 million population.

[Insert Figures 4 and 5 about here]

Next, we zoom in and plot the daily estimates of Equation (1) in event study-type graphs. Figure 4a shows *all cause admissions* per 100,000 population and Figure 4b *cardiovascular admissions* per 100,000 population. Despite conservative two-way clustering, we are able to identify even daily effects in a very precise manner. Please note that the event study graphs do *not* represent simple descriptive graphs but compare the effects in the treatment group relative to the control group (Figure 1) after having netted out of seasonal and weekday confounders as formalized by Equation (1).

The two event study graphs show a characteristic four-day pattern of decreases in admissions: We observe significant decreases in overall and cardiovascular admissions on days one to four after the time shift. The effect is strongest on the Monday after the clocks are set back, and it decreases smoothly over the next three days before it disappears on day five. The decrease for cardiovascular admissions equals about 10 avoided admission per one million population for four days.

In robustness checks, we obtain exactly the same pattern using the full sample (Figure A1, Appendix) as well as *heart attacks* and *injuries* (Figure 5). The consistency of these patterns for even heart attacks is reassuring.⁶

Finally, we examine hospital admissions due to *drug overdosing*, which arguably has a weaker theoretical link with sleep. Illicit drugs are highly addictive, which limits the extent to which additional sleep can help prevent those who are on the margin of overdose from being hospitalized. As such, we do not expect to see a strong effect on drug-related hospital admissions. Indeed, Figure 6 does not show much of an effect.

[Insert Figure 6 about here]

In conclusion, we interpret the similarity of these four-day patterns as strong support for our identification strategy. The implication is that additional sleep leads to immediate health improvements for people who are *on the margin* of being hospitalized and prevents about ten heart admissions per one million population for four days. (On average per day, 7815 people are admitted to German hospitals because of heart issues, cf. Table A1.) This finding is very consistent with, and underscores, the medical advice that people on the margin of having acute heart failure should get sufficient bed rest (Millane et al. 2000).

4.3 Alternative Mechanisms

Next, we explore alternative channels through which the DST transition may affect hospital admissions. One possible channel is through a shift in ambient light. As the clocks “fall back” by one hour, sunrise and sunset both occur at earlier times. One could hypothesize that, because

⁶ Note that the German data do not allow us to distinguish between emergency room visits, elective visits and other type of admission. We solely see the primary diagnosis and know that the patient stayed overnight, which excludes ambulatory elective surgeries.

mornings get brighter earlier, people are more likely to exercise in the morning following the transition (and less likely to exercise in the evening). To test for such an exercise effect, we use a question in the BRFSS on exercising and estimate our standard model in Equation (1). Figure 7a shows the daily effects. In line with Giuntella and Mazzonna (2019), we find no evidence that the frequency of exercise changes due to the time shift.

[Insert Figure 7 about here]

Next, we stratify the effects by weather conditions using the German Hospital Census. We use data from more than one thousand ambient weather monitors on a daily basis from 2000 to 2008. The underlying hypothesis is that weather conditions determine how and where individuals spend their time (Gebhart and Noland, 2014); better outdoor conditions should also indicate whether changes in exercising behavior play a confounding role. Table A2 stratifies the effects by (i) temperature, (ii) rainfall, (iii) sunshine, and (iv) cloudiness. Methodologically, we run our standard model, control for weather conditions and interact DST_{id} with the weather measures in the column headers. Consistent with the absence of changes in exercising (Figure 7a), there is no evidence that ambient conditions matter. None of the interaction terms between the four weather measures and DST_{id} is statistically significant.

A shift in ambient light can also affect traffic accidents (Hicks et al., 1998; Smith, 2016). This could potentially explain the significant reduction in admissions due to injuries. However, traffic accidents would not be able to explain why we observe reductions in admissions across many disease categories that are unrelated to accidents, such as admissions for cardiovascular diseases.

Another potential confounding factor is crime. Doleac and Sanders (2015) show that robberies decrease in the days following the DST transition in spring (when evenings get dark later). They find no significant effects on crime rates in fall. If there was a significant robbery effect, robberies

would likely *increase* following the time shift in the fall (because it gets dark sooner), and thus have adverse health effects, opposite what we find. It is also unlikely to explain health benefits across a wide range of disease categories.⁷

The fall DST transition increases the length of the Sunday from 24 to 25 hours. This may affect hospital admissions (or survey responses) in ways unrelated to sleep. However, because the day is longer, it will result in *more* admissions, opposite our findings. This mechanism also cannot explain the persistent health effects that we find over four days.

Finally, we estimate placebo regressions. Our first placebo test, using BRFSS, is having received a flu shot in the *past year* as an outcome measure. This outcome is, by construction, unrelated to getting additional sleep. As expected, Figure 7b shows no impact on this outcome.

[Insert Figure 8 about here]

Our second placebo test uses the hospital data to conduct the following permutation test: We start in July of each year and select six-week windows of data as illustrated in Figure 1. Then, we run our standard model with aggregated effects at the weekly level, pretending that the fourth week was the week of the time shift. Next, we move the six-week window one week further into August and repeat the approach. We permute until week six of our selected sample hits the true week of the time shift and continue with six-week windows until end of the year.⁸ As such, we obtain 23

⁷ While both effect sizes—on robberies and fatalities—are cleanly identified by the studies just cited, they are rather small and unlikely to confound our population health estimates. According to Doleac and Sanders (2015), in spring, the number of avoided robberies decrease by about 2 per 10 million people. Smith (2016) finds that the spring change leads to 30 more deaths for the entire U.S. These numbers certainly would not bias the survey estimates for the U.

S. As for the hospital admission data, our “Injury Admissions per 1 Million Population” outcome category should capture these effects.

⁸ The true DST week is never included in these placebo six week samples.

weekly placebo estimates. Figure 8 plots the distribution of these weekly placebo estimates along with the true estimate. Clearly, the decrease in admissions following the time shift does not fall within the statistical placebo estimate distribution.

4.5 Quantifying the Economic Benefits of One Additional Hour of Sleep

When considering policies to tackle the public health issue of sleep-deprivation, it is important to quantify the potential benefits of encouraging people to sleep more. In this section, we monetize the economic benefits of avoided hospital admissions. We also estimate other benefits based on studies that show improved work productivity (Gibson and Schrader, 2018) and avoided traffic fatalities (Smith, 2016). These calculations are based on several assumptions, but provide a basic framework for such an exercise.

[Insert Table 3 about here]

Table 3 shows the estimates. We first monetize the value of avoided hospital admissions from the end of DST in the fall. Figure 4a implies 100 fewer admissions per 1 million population over four days. Columns (1) to (3) in Table 3 show that the benefits of avoided hospital stays can be decomposed into a per person €2,000 for medical costs, €450 for lost labor as well as €550 for lost quality of life during hospital stays.

Next, we assess the value of increases in work productivity when sleep-deprived employees gain more sleep. According to Gibson and Schrader (2018), the short-term wage returns for an additional hour of sleep equals 1.1% of the wage. Given the average daily wage of \$230 in the U.S., this translates into \$10 over four days. Assuming that these gains only apply to the ten percent chronically sleep deprived full-time employed Americans (Knutson et al. 2010), it would sum to \$500 thousand per 1 million population (column (4), Table 3).

Finally, Smith (2016) quantifies the number of avoided traffic fatalities with 30 for the entire U.S. (0.09 per 1 million population). Evaluated at \$5 million per life saved (Kniesner et al. 2010), we obtain values for saved statistical lives of around \$450 thousand per 1 million population (column (5), Table 3).

In conclusion, we estimate that the welfare benefits of a nighttime extension by an hour sum to about \$1.3 million per 1 million population.

5. CONCLUSION

This paper exploits the quasi-experimental nature of Daylight Saving Time (DST) to assess the health benefits of increasing people's sleep. We find that people sleep significantly more in the short-run when they gain an additional hour at night following the DST "fall back." Moreover, the share of people who unintentionally fall asleep during the day drops significantly for four days. In addition, we find that hospital admissions drop sharply for four days as well. For example, cardiovascular admissions decrease by ten per one million population. We find no effect for placebo outcomes, which have weaker or no theoretical links to sleep, such as drug overdosing or having received a flu shot.

Because exogenous shifters of sleep are very rare in real world settings, our study is one of very few causal studies on the health benefits of sleep (one of the exemptions is Giuntella and Mazzonna, 2019). To identify effects, we use a large survey from the U.S. and the census of hospital admissions from Germany. Properly investigating the impact of the nighttime extension on health *on a daily level* requires powerful and representative data. These are crucial to estimate rich econometric specifications that consider weekday effects in addition to general and specific seasonal adjusters.

Our findings have important implications for public policy. Sleep deprivation is becoming a widespread problem in many developed countries—the CDC has recently declared it a “public health epidemic” (CDC 2014). Almost a third of Americans report sleeping six or fewer hours, significantly less than the CDC-recommended minimum of 7 hours (cf. Sheehan et al. 2019). The findings from our study reinforce the need to devise policies to reduce sleep deprivation in the population.

The evidence in this paper is also bolstered by other recent economic studies that identify work productivity effects as a result of more sleep (Gibson and Schrader, 2018), decreases in obesity (Giuntella and Mazzonna, 2019), better cognitive skills (Giuntella et al. 2017) or fewer traffic fatalities (Smith, 2016). In the last part of the paper, we attempt to categorize, standardize, and monetize the various benefits that this paper and companion research in economics identifies. Under some assumptions, we assess the total societal benefits of gaining one hour of sleep with about \$1.3 million per 1 million population. The benefits can be decomposed into work productivity, hospitalization, and mortality effects.

The main objective of this paper is to provide evidence of a causal relationship between Daylight Saving Time transitions, sleep and health. We do not intend to draw conclusions about the overall welfare effects of Daylight Saving Time. We also would like to point to a caveat: our reduced-form approach is well-suited for the identification of causal and immediate intent-to-treat effects, but less suited to identify long-term effects of sleep. Based on sleep habits, sleep may affect mood, cognitive skills and health cumulatively over time in the long-run. Alternatively, it is possible that the human body is able to adapt to (adverse) sleeping conditions. Field experiments have the power to find answers to these questions (cf. Tepeđino et al. 2017). More

research is necessary to better understand how improvements in sleep quality may improve living quality, education and labor market outcomes as well as life expectancy.

LITERATURE

Almond, D., & Currie, J. (2011): Killing Me Softly: The Fetal Origins Hypothesis, *Journal of Economic Perspectives*, 25(3): 153-172.

Banks, S., Dinges, D. (2007): Behavioral and Physiological Consequences of Sleep Restriction. *Journal of Clinical Sleep Medicine*, 3(5): 519-28.

Barnes, C. M., Wagner, D. T. (2009): Changing to Daylight Saving Time Cuts into Sleep and Increases Workplace Injuries, *Journal of Applied Psychology*, 94 (5): 1305-1317.

Becker, G. S. (1964). *Human Capital: A Theoretical and Empirical Analysis*, with Special Reference to Education, 1st edition, Chicago, University of Chicago Press.

Berk, M., Dodd, S., Hallam, K., Berk, L., Gleeson, J., & Henry, M. (2008): Small Shifts in Diurnal Rhythms are Associated with an Increase in Suicide: The Effect of Daylight Saving, Sleep and Biological Rhythms, 6(1): 22-25.

Biddle, J. E., & Hamermesh, D. S. (1990): Sleep and the Allocation of Time, *Journal of Political Economy*, 98(5): 922-943.

Billari, F., Giuntella, O., & Stella, L. (2018): Broadband Internet, Digital Temptations, and Sleep, *Journal of Economic Behavior and Organization*, 153: 58-76.

Cameron, A. C., Gelbach, J. B., & Miller, D. L. (2011): Robust Inference With Multiway Clustering, *Journal of Business & Economic Statistics*, 29(2): 238-249.

Carrell, S. E., Maghakian, T., West, J. E. (2011): A's from Zzzz's? The Causal Effect of School Start Time on the Academic Achievement of Adolescents, *American Economic Journal: Economic Policy*, 3(3): 62-81.

Carroll, L. (2016): It's Daylight Saving Time! 6 Tips to Help You Deal with the Change, March 11, 2016, <https://www.today.com/health/ready-time-change-why-you-should-grab-banana-avoid-blue-t79346>, April 23, 2019.

Cawley, J. & Ruhm, C. J. (2011): "The Economics of Risky Health Behaviors," in Eds: Pauly, M. V.; McGuire, T. G., & Barros, P. P.: *Handbook of Health Economics*, Volume 2, Chapter Three - Pages 95-199. Elsevier.

CDC (2014): Insufficient Sleep Is a Public Health Epidemic, <http://www.accqsleeplabs.com/wp-content/uploads/2014/10/CDC-report.pdf>, retrieved October 19, 2018.

- Cunha, F., & Heckman, J. (2007): The Technology of Skill Formation, *American Economic Review*, 97(2): 31-47.
- Currie, J., Graff Zivin, J., Mullins, J., & Neidell, M. (2014): What Do We Know About Short- and Long-Term Effects of Early-Life Exposure to Pollution? *Annual Review of Resource Economics*, 6(1): 217-247.
- Doleac, J. L., & Sanders, Nicholas. J. (2015): Under the Cover of Darkness: How Ambient Light Influences Criminal Activity, *Review of Economics and Statistics*, 97(5): 1093-1103.
- Ford, Earl S., Timothy J. Cunningham, and Janet B. Croft (2015): Trends in self-reported sleep duration among US adults from 1985 to 2012, *Sleep* 38(5): 829-832.
- Gebhart, K., & Noland, R. (2014): The Impact of Weather Conditions on Bikeshare Trips in Washington, DC, *Transportation*, 41(6): 1205-1225.
- German Federal Statistical Office (2017): *Statistical Yearbook 2017 for the Federal Republic of Germany*. Metzler-Poeschel.
- Gibson, M., & Shrader, J. (2018): Time Use and Labor Productivity: The Returns to Sleep, *Review of Economics and Statistics*, 100(5): 783-798.
- Giuntella, O., & Mazzonna, F. (2019): Sunset Time and the Effects of Social Jetlag. Evidence from US Time Zone Borders, *Journal of Health Economics*, forthcoming.
- Giuntella, O., Han, W. & Mazzonna, F. (2017): Circadian Rhythms, Sleep and Cognitive Skills: Evidence from an Unsleeping Giant, *Demography*, 54(5): 1715–1742.
- Graff Zivin, J., & Neidell, M. (2013): Environment, Health, and Human Capital, *Journal of Economic Literature*, 51(3): 689-730.
- Grossman, M. (1972): On the Concept of Health Capital and the Demand for Health, *Journal of Political Economy*, 80(2): 223-255.
- Hamermesh, D. S., Myers, C. K., & Pockock, M. L. (2008): Cues for Timing and Coordination: Latitude, Letterman, and Longitude, *Journal of Labor Economics*, 26(2): 223-246.
- Hanigan, I., Hall, G., & Dear, K. B. G. (2006): A Comparison of Methods for Calculating Population Exposure Estimates of Daily Weather for Health Research, *International Journal of Health Geographics*, 38(5): 1.
- Harrison, Yvonne. (2013): The impact of daylight saving time on sleep and related behaviours, *Sleep medicine reviews*, 17(4): 285-292.
- Hausman, C. & Rapson, D. S. (2018): Regression Discontinuity in Time: Considerations for Empirical Applications, *Annual Review of Resource Economics*, 10:1

- Hillman, D. R., Murphy, A. S., Antic, R., & Pezzullo, L. (2006): The Economic Cost of Sleep Disorders, *Sleep*, 29(3): 299-305.
- Hicks, G. J., Davis, J. W., & Hicks, R. A. (1998): Fatal Alcohol-Related Traffic Crashes Increase Subsequent to Changes to and from Daylight Savings Time, Perceptual and Motor Skills, 86(3): 879-882.
- Janszky, I., Ahnve, S., Ljung, R., Mukamal, K. J., Gautam, S., Wallentin, L., & Stenestrand, U. (2012): Daylight Saving Time Shifts and Incidence of Acute Myocardial Infarction—Swedish Register of Information and Knowledge About Swedish Heart Intensive Care Admissions (RIKS-HIA), *Sleep Medicine*, 13(3): 237-242.
- Jiddou, M. R., Pica, M., Boura, J., Qu, L., & Franklin, B. A. (2013): Incidence of Myocardial Infarction With Shifts to and From Daylight Savings Time, *The American Journal of Cardiology*, 111(5): 631-635.
- Jin, L. & Ziebarth, N. R. (2016): Sleep and Human Capital: Evidence from Daylight Saving Time, Canadian Centre for Health Economics, Working Papers 160001. <https://ideas.repec.org/p/cch/wpaper/160001.html>, retrieved on April 23, 2019.
- Kesternich, I., Siflinger, B., Smith, J., & Winter, J. (2015): Individual behavior as a pathway between early-life shocks and adult health: Evidence from hunger episodes in post-war Germany, *Economic Journal*, 125, F372-F393.
- Killgore, W. D. S. (2010): Effects of Sleep Deprivation on Cognition, in Kerkhof, G. A. & Van Dongen H. P. A.: *Progress in Brain Research*, Vol. 185: pages 105-129.
- Kniesner, T., W. Viscusi, and J. Ziliak (2010): Policy Relevant Heterogeneity in the Value of Statistical Life: New Evidence from Panel Data Quantile Regressions, *Journal of Risk and Uncertainty* 40 (1): 15-31.
- Kotchen, M. J., & Grant, L. E. (2011): Does Daylight Saving Time Save Energy? Evidence from a Natural Experiment in Indiana, *The Review of Economics and Statistics*, 93(4): 1172-1185.
- Kountouris, Y., & Remoundou, K. (2014): About Time: Daylight Saving Time Transition and Individual Well-Being, *Economics Letters*, 122(1): 100-103.
- Kuehnle, D., & Wunder, C. (2016): Using the Life Satisfaction Approach to Value Daylight Savings Time Transitions: Evidence from Britain and Germany, *Journal of Happiness Studies*, 17(6): 2293-2323.
- Lahti, T., Leppämäki, S., Lönnqvist, J., Partonen, T. (2008): Transitions Into and Out of Daylight Saving Time Compromise Sleep and the Rest-Activity Cycles, *BMC Physiology*, 8(1), 1.

- Liu Y., Wheaton A.G., Chapman D.P., Cunningham T.J., Lu H., Croft J.B. (2016): Prevalence of Healthy Sleep Duration among Adults 2014— United States, MMWR Morbidity and Mortality Weekly Report, 65:137–141.
- Millane, T., Jackson, G., Gibbs, C. R., & Lip, G. Y. H. (2000). Acute and chronic management strategies. *BMJ : British Medical Journal*, 320(7234), 559–562.
- Moore, P. J., Adler, N. E., Williams, D. R., & Jackson, J. S. (2002): Socioeconomic Status and Health: The Role of Sleep, *Psychosomatic Medicine*, 64(2): 337-344.
- Mullainathan, S. (2014): Get Some Sleep, and Wake Up the GDP, *New York Times*, February 2, 2014 http://www.nytimes.com/2014/02/02/business/get-some-sleep-and-wake-up-the-gdp.html?_r=0
- Mullington, J. M., Haack, M., Toth, M., Serrador, J. M., & Meier-Ewert, H. K. (2009): Cardiovascular, Inflammatory, and Metabolic Consequences of Sleep Deprivation, *Progress in Cardiovascular Diseases*, 51(4): 294-302.
- OECD (2017). OCED Statistics, <http://stats.oecd.org>, retrieved November 20, 2017.
- Sexton, A. L., & Beatty, T. K. M. (2014): Behavioral Responses to Daylight Savings Time, *Journal of Economic Behavior & Organization*, 107(PA): 290-307.
- Sheehan, C. M., Frochen, S. E., Walsemann, K M., & Ailshire, Jennifer A. (2019): Are U.S. adults Reporting Less Sleep?: Findings from Sleep Duration Trends in the National Health Interview Survey, 2004–2017, *Sleep*, 42(2): zsy221.
- Smith, A. C. (2016): Spring Forward at Your Own Risk: Daylight Saving Time and Fatal Vehicle Crashes, *American Economic Journal: Applied Economics*, 8(2): 65-91.
- Spaeth, A. M., Dinges, D., & Goel, N. (2013): Effects of Experimental Sleep Restriction on Weight Gain, Caloric Intake, and Meal Timing in Healthy Adults. *Sleep*. 36. 981-990.
- Taheri, S., Lin, L., Austin, D., Young, T., & Mignot, E. (2004): Short Sleep Duration Is Associated with Reduced Leptin, Elevated Ghrelin, and Increased Body Mass Index, *PLoS Medicine*, 1(3): e62.
- Tepedino, P. B., Rao, G., Schilbach, F., Schofield, H. & Toma, M. (2017): Pre-analysis Plan Outline for ‘Sleepless in Chennai: The Consequences of Sleep Deprivation among the Urban Poor’, <https://www.socialscienceregistry.org/docs/analysisplan/1365/document>, retrieved May 10, 2018.
- U.S. Census Bureau (2012). US Census 2010. <http://2010.census.gov/2010census/data>, retrieved November 20, 2017.

Van Dongen, H. P. A., Maislin, G., Mullington, J. M., & Dinges, D. F. (2003): The Cumulative Cost of Additional Wakefulness: Dose-Response Effects on Neurobehavioral Functions and Sleep Physiology from Chronic Sleep Restriction and Total Sleep Deprivation, *Sleep*, 26(2): 117-126.

Van Hare, H. (2019): Your Daylight Saving Time Survival Guide: Spring Forward Painlessly with this Comprehensive Guide, March 1, 2019, <https://www.thedailymeal.com/healthy-eating/daylight-saving-survival-guide>, retrieved on April 23, 2019.

Witte, D. R., Grobbee, D.E., Bots, M.L. & Hoes, A.W. (2005): A Meta-Analysis of Excess Cardiac Mortality on Monday, *European Journal of Epidemiology*, 20(5): 401-406.

Figures and Tables

Figure 1: Sample Selection of Main Models—Extracting 6 Weeks around DST Transition






Figure 2: Nonparametric Plot of Sleep Duration in DST Week vs. Before and After




Source: BRFSS, 2013-2016. The gray dashed line plots average sleep duration on the week day level in the weeks before and after the transition. The solid blue line plots sleep duration for the seven days around the transition in fall. The transition occurs Sunday at 2am, represented by the black vertical line.

Figure 3a,b: Effects of Nighttime Extension on Sleep and Unintentionally Falling Asleep




Source: BRFSS, 2013-2016 (Panel a) and 2009-2010 (Panel b). Equation (1) is estimated and daily effects plotted.

Figure 4a,b: Effects of Nighttime Extension on Total and Cardiovascular Hospital Admissions




Source: German Hospital Census, 2000-2008. Equation (1) is estimated and daily effects plotted.

Figure 5: Effects of Nighttime Extension on Heart Attacks and Injuries




Source: German Hospital Census, 2000-2008. Equation (1) is estimated and daily effects plotted.

Figure 6: Effects of Nighttime Extension on Drug Overdosing




Source: German Hospital Census, 2000-2008. Equation (1) is estimated and daily effects plotted.

Figure 7a, b: Placebo Tests—Effects on Exercising and Flu Shot in the Past Year



Source: BRFSS. Equation (1) is estimated and daily effects plotted.

Figure 8: Permutation Test Comparing Placebo Effects to Fall DST Transition Week



Source: German Hospital Census, 2000-2008. Equation (1) is estimated and placebo effects plotted for six-week windows in the second half of the year. Red line indicates true fall DST weekly coefficient estimate.

Table 1: The Effects of Fall DST on Sleep Duration and Tiredness

	(1) Hours of Sleep	(2) Unintentionally fell asleep during day at least once in past 30 days
Week of Transition (End of DST)	0.026*** (0.010)	-0.044** (0.022)
Controls		
State FE	X	X
Halloween	X	X
Day of Week * Month FE	X	X
Month * Year FE	X	X
Linear & quad. time trend	X	X
<i>Mean of dep. Var.</i>	7.06	0.35
Observations	174,503	10,833

Notes: * p<0.1, ** p<0.05, *** p<0.01. The data are from BRFSS. Standard errors in parentheses are clustered at the date level. Regressions are probability-weighted. *Week of Transition* is an indicator that equals 1 if the interview is on the Sunday of DST transition or one of the subsequent six days. The column headers describe the dependent variables used in each column. Each column is one model as in Equation (1). The sample period for column (1) is 2013-2016. The sample period for column (2) is 2009-2010, as the survey question was experimented by several states at the time and discontinued since. This includes six states (Georgia, Hawaii, Illinois, Louisiana, Minnesota, and Wyoming) in 2009 and nine states in 2010 (Arkansas, Connecticut, Delaware, District of Columbia, Hawaii, Minnesota, Missouri, Nevada, and Oregon).

Table 2: The Effects of Fall DST on Hospitalizations by Disease Type

	All cause admission rate (1)	Cardiovascular admission Rate (2)	Heart attack rate (3)	Injury admission rate (4)	Metabolic adm. rate (5)	Suicide attempt rate (7)	Drug Overdosing (8)
Week of Transition (End of DST)	-4.9556*** (1.1139)	-0.7195*** (0.1589)	-0.0882*** (0.02611)	-2.7121*** (0.6869)	-0.1874*** (0.0385)	-0.0276** (0.0128)	-0.0044 (0.0055)
Controls							
County FE	X	X	X	X	X	X	X
Vacation & holiday FE	X	X	X	X	X	X	X
Day of Week * Month FE	X	X	X	X	X	X	X
Month*Year Fixed Effects	X	X	X	X	X	X	X
Linear & quadr. time trend	X	X	X	X	X	X	X
Socioeconomic covariates	X	X	X	X	X	X	X
<i>Mean of dep. variable</i>	59.77	9.53	1.59	57.56	0.32	0.09	0.32
R ²	0.8469	0.5675	0.1510	0.2067	0.3095	0.0179	0.0008
Observations	336,604	336,604	336,604	336,604	336,604	336,604	336,604


Note: * p<0.1, ** p<0.05, *** p<0.01. Standard errors are in parentheses and two-way clustered at the county and date level. *Week of Transition* is an indicator variable that equals 1 if the interview date is on the DST Sunday or one of the following six days. Table A1 lists the dependent variables for as displayed in the column header. Each column is one model as in Equation (1). All admission rates are per 100,000 except for *Injuries*, *Suicides* and *Drug Overdosing* (per 1,000,000).

Table 3: Decomposing and Monetizing Benefits of Additional Sleep

	<i>Health Effects</i>			<i>Productivity Effects</i>	<i>Mortality Effects</i>
	German Hospital Census (Fig 4a; Table 2)			Gibson and Schrader (2018)	Smith (2016)
	Healthcare Costs	Labor Productivity	QALYs	Increase in Work Productivity	Avoided Deaths
	€500 per day *4 days	€150 per day *4 days	(\$100K/365) *0.5 *4 days	+1.1% at \$230 daily wage *4 days	30 fatalities in U.S. (0.09 per 1M pop.)
Benefit for individual	=€2000 *100	=€450 *(100/3)	=€550 *100	= \$10 *10% sleep deprived employees: (0.1*161M)/320M	*\$5M per VSL
Per 1M pop.	=€200K	=€15K	= €55K	= \$500K	= \$450K


Appendix A

Figure A1: The Effects of Fall DST Transition on Total Admissions, Full Sample



Source: German Hospital Census, 2000-2008. Equation (1) is estimated and daily effects plotted.

Figure A1: Effects of Spring DST on Injury Admissions



Source: German Hospital Census, 2000-2008. Equation (1) is estimated for spring DST and daily effects plotted.

Table A1: Descriptive Statistics

	<i>Mean</i>	<i>Std.Dev</i>	<i>Min.</i>	<i>Max.</i>	<i>Obs.</i>
BRFSS					
Hours of sleep	7.06	1.46	1	24	174,503
Unintentionally fell asleep at least once in past 30 days	0.346	0.476	0	1	10,833
German Hospital Census					
Dependent Variables					
Total admission rate per 100,000	59.7681	25.7333	N/A	N/A	336,604
Cardiovascular admission rate per 100,000	9.5339	4.9525	N/A	N/A	336,604
Heart attack admission rate per 100,000	1.5909	1.4035	N/A	N/A	336,604
Injury admission rate per 1 million	56.5571	26.6603	N/A	N/A	336,604
Suicide attempt rate per 1 million	0.3219	1.6754	N/A	N/A	336,604
Drug overdosing rate per 1 million	0.0892	0.8594	N/A	N/A	336,604
Socio-Demographic Individual Controls					
Female	0.5420	0.0671	0	1	336,604
Surgery needed	0.3715	0.1478	0	1	336,604
Died in hospital	0.0249	0.0230	0	0.5	336,604
Private hospital	0.1177	0.1813	0	1	336,604
Age Group 0-2 years	0.0619	0.0416	0	0.5556	336,604
....	336,604
Age Group 65-74 years	0.0161	0.0182	0	0.3333	336,604
>74 years	0.0034	0.0082	0	0.5	336,604
Annual County-Level Controls					
Hospital per county	4.8196	5.4690	0	76	336,604
Hospital beds per 10,000	1204.02	1574.54	0	24,170	336,604
Unemployment rate in county	10.37	5.29	1.6	29.3	336,604
Physicians per 10,000	153.96	53.18	69	394	336,604
GPD per resident (in Euro)	25,235	10,219	11,282	86,728	336,604
Seasonal Controls					
Fall Vacation	0.0977	0.2969	0	1	336,604
Week End DST	0.0862	0.2807	0	1	336,604

Source: Sleep variables are obtained from the Behavioral Risk Factor Surveillance System (BRFSS). *Hours of sleep* is obtained from 2013-2016, and *unintentionally fallen asleep* is obtained from 2009-2010 due to limited data availability. The hospital admission data are from the German Hospital Census 2000-2008, Federal Institute for Research on Building, Urban Affairs and Spatial Development (2012). The hospital admission data are aggregated at the county-day level and normalized per 100,000 population. Note that both nominator and denominator refer to the county of residence. The data excludes military hospitals and hospitals in prisons. Note that German data protection laws prohibit us from reporting min. and max. values. The socio-demographic individual controls are also aggregated at the county-day level. The seasonal controls only vary between days, not across counties. The annual county-level controls vary between the counties and over years, but not within years. Between 2000 and 2008, Germany had up to 468 different counties. Mostly, due to mergers and reforms of the administrative boundaries, the number of counties varies across years.

Linking Hospital with Official Weather Data

Weather Data. The weather data are provided by the German Meteorological Service (*Deutscher Wetterdienst (DWD)*). The DWD is a publicly funded federal institution and collects information from hundreds of ambient weather stations which are distributed all over Germany. Daily information on the average temperature, rainfall, hours of sunshine and cloudiness from up to 1,044 monitors and the years 2000 to 2008 are used.

We extrapolate the point measures into space using inverse distance weighting. This means that the measures for every county and day are the inverse distance weighted average of all ambient monitors within a radius of 60 km (37.5 miles) of the county centroid (Hanigan et al. 2006).

Socioeconomic Background Data. Because the *Hospital Admission Census* only contains gender and age, we link yearly county-level data with the hospital data. We merge in county-level information on *GDP per resident*, the *unemployment rate*, the *number of physicians per 10,000 pop.*, the *number of hospitals in county* as well as the *number of hospital beds per 10,000 pop*

Table A2: Effects of Fall DST Transition on Admissions by Weather Conditions

	All cause admission rate			
	(1)	(2)	(3)	(4)
	Temp.	Rainfall	Sunshine	Cloudiness
DST * [column header]	-0.2378 (0.2329)	0.0991 (0.1396)	-0.2095 (0.3431)	0.4458 (0.5083)
DST (3am → 2am in fall)	-3.1330* (1.8671)	-5.1829*** (1.2059)	-4.4812*** (1.1962)	-7.6059** (3.3876)
Controls				
Halloween, Vacation FE	X	X	X	X
Day of Week * Month FE	X	X	X	X
Month * Year FE	X	X	X	X
Linear & quadratic trend	X	X	X	X
Socioecon. covariates	X	X	X	X
Weather and pollution controls	X	X	X	X
R ²	0.8372	0.8372	0.8373	0.8373
Observations	336,604	336,604	336,604	336,604

Notes: *** Significant at 1% level, ** 5%, * 10%. Standard errors in parentheses are two-way clustered at the date and county level. *DST* are indicator variables equal to 1 if the interview is on the DST Sunday or one of the following 6 days. The dependent variable is the all cause hospital admission rate per 100,000 pop. at the daily county level. Appendix A describes the weather measures and how they are linked to the Hospital Census on a daily county-level basis. Each column is one model as in Equation (1).

Appendix B: Measurement of Outcome Variables

This paper uses self-reported measures on sleep and tiredness from the BRFSS as well as administrative hospital admission data from Germany. Together these represent a broad set of measures from different countries to validate our findings.

First, the BRFSS tiredness measure refers to “in the last 30 days”, which may introduce measurement error and a non-straightforward interpretation when used as outcome. Assume that there was no recall bias or measurement error and everybody would provide accurate answers. Further, assume that DST would affect respondents for four days. Then, those interviewed on the day of the DST transition would report their average sleep duration for 29 days prior to DST and the first post-DST day; those interviewed on Monday would report their sleep for two post-DST days, and so on. Because our standard approach assigns respondents in weeks $t+2$ and $t+3$ to the control group status (Figure 1), our estimates would be downward biased as the retrospective 30-day responses would be affected by DST as well. In practice, however, we expect recall biases and that respondents overweigh days closer to the interview day. In robustness checks, we assign respondents in weeks $t+2$ and $t+3$ to the treatment group and the results hold up.

Second, we use administrative hospitalization data: German geography, combined with the institutional setting of the German health care system, makes it very plausible that variations in hospitalizations represent serious population health effects. Germany has 82 million residents living in an area, which has roughly the size of the U.S. state Montana. Thus, the average German population density is seven times higher than the U.S. population density and 231 vs. 32 people per km^2 (U.S. Census Bureau, 2012; German Federal Statistical Office, 2017). The hospital bed density is also much higher. Per 100,000 population, Germany has 824 hospital beds, while the U.S. has 304 beds (OECD, 2017). Hence, geographic hospital access barriers, such as travel distances, are low in Germany. Moreover, the German uninsurance rate is below 0.5%. The public

health care system covers 90% of the population and copayment rates in the public scheme are uniform and low. The overwhelming majority of hospitals can be accessed independently of insurance status and free choice of providers exist (no provider networks).